

## Role of different fertilization on growth, yield and quality evaluation in maize (*Zea mays* L.) and cowpea (*Vigna unguiculata* L.) intercropping

ANISHARANI<sup>1</sup>, ARSHVEER SINGH<sup>2</sup>, GURPREET SINGH<sup>3</sup> AND PRADEEP KUMAR SRIVASTAVA

Associate Professor, Department of Agriculture, Khalsa College, Patiala-147001

EmailID: drpksoil@gmail.com

### Abstract

A field experiment was conducted during Zaid season 2022-2023. The experiment was conducted in split plot design and replicated thrice. The treatment in main plot consisted of three intercropping and in sub plots five different organic manures and different levels of fertilizers. Soil of the experiment field was clayey in texture had soil slightly alkaline, medium organic carbon (0.71%), low in available nitrogen (246.64 kg ha<sup>-1</sup>), medium in available phosphorous (21.6 kg ha<sup>-1</sup>) and high in available potassium (282 kg ha<sup>-1</sup>). A keen observation of data revealed that application of organic manures and different levels of fertilizers with intercropping both significantly influenced the growth parameters, yield parameters and yield, economic attributes like gross return, net return and B:C ratio was significantly maximum at IC<sub>2</sub> maize + cowpea (2:1) with different organic manures and levels of fertilizers application F<sub>5</sub> (50% RDF + 5 t FYM ha<sup>-1</sup> + 2 t VC ha<sup>-1</sup> + 2 t PM ha<sup>-1</sup>).

Keywords: (*Zea mays* L.), (*Vigna unguiculata* L.), intercropping, organic manures

### Introduction

Maize (*Zea mays* L.) is one of the most important cereal crops in the worldwide. It belongs to family Gramineae and originated from the Central America and Mexico. It belongs to C<sub>4</sub> plant. Maize is highly valued cereals in the world that is why it is called "Queen of cereals". Having haploid number of 10 chromosomes (2n=20). In India, the average area is 9.76 million hectare with production of 26.14 million tones and having productivity 2629.28 kg ha<sup>-1</sup> (Mishra *et al.* 2019). Maize crop absorbs large quantity of nutrients from soil for complete their life cycle. (Mahmood *et al.* 2017) adopted the different management practices to increase the maize yield

and their production. Use of organic manures to increase the soil structure, soil texture and water holding capacity in the soil is improved. The application of UREA and DAP is help to increase the yield of maize.

Cowpea (*Vigna unguiculata* L.) known as "lobia" is used as a pulse crop, a fodder crop and green manure crop. It is a legume crop. It is also known as black eye pea (Kaviraj *et al.* 2017). It is rich in protein that's why it is also known as vegetable meat. It is used for both men as well as animal feed. Cowpea is mainly grown in Africa. About 90% of the total world acreage is in Africa.

<sup>1</sup>Research Scholar, Department of Agriculture, Khalsa College, Patiala-147001

<sup>2</sup>Assistant Professor, Shaheed Bhagat Singh State University, Ferozpur-152004

<sup>3</sup>Assistant Professor, Department of Agriculture, Khalsa College, Patiala-147001

In India the area under cowpea is 3.9 million hectares with annual production of 2.21 million tones with a national productivity of 638 kg ha<sup>-1</sup>.

Intercropping of legumes and cereals is an old practice in tropical agriculture that dates back to ancient civilization. Cereal and Legume intercropping is more productive and profitable cropping system in comparison with solitary cropping (Evans *et al.* 2001). The main objective of intercropping has been to maximize use of resources such as space, light and nutrient (Li *et al.* 2003). The importance of intercropping in farming practices has long been recognized in India (Reddy *et al.* 2004).

### Materials and Methods

The present investigation was carried out during *Zaid* season of the year 2022-2023 at Agronomy Research Farm Dhablan (Patiala). The experimental site is situated at about 30°19' North Latitude and 76°24' East Longitude at an altitude of about 250 m above the mean sea level. It is located in south eastern direction in Punjab state and North West India. The experiment site falls in Indo-Gangetic plains. During growing season (February to June), the weekly maximum and minimum temperature ranged from 24.7 °C – 44.4 °C and 10.4 °C – 27.7 °C, respectively. The average relative humidity ranged 90.2% to 43.0%. The total rainfall received during the crop period was 17.3 mm. The soil of experiment site was clayey having 0.71% organic carbon, 246.64 kg ha<sup>-1</sup>, 21.6 kg ha<sup>-1</sup> and 282 kg ha<sup>-1</sup> of available N, P<sub>2</sub>O<sub>5</sub>, K<sub>2</sub>O respectively. The experiment was laid out in a SPD (split plot design) and replicated thrice. The main plot having three intercropping and the sub plot having five organic manures and different levels of fertilizers. The intercropping IC<sub>1</sub> (5:1), IC<sub>2</sub> (2:1), IC<sub>3</sub> (3:1). The fertilizer levels F<sub>1</sub>: 100% RDF, F<sub>2</sub>: 75% RDF + 10 t FYM ha<sup>-1</sup>, F<sub>3</sub>: 50% RDF + 5 t FYM ha<sup>-1</sup> + 2 t VC ha<sup>-1</sup>, F<sub>4</sub>: 50% RDF + 5 t FYM ha<sup>-1</sup> + 2 t PM ha<sup>-1</sup>, F<sub>5</sub>: 50% RDF + 5 t FYM ha<sup>-1</sup> + 2 t VC ha<sup>-1</sup> + 2 t PM ha<sup>-1</sup>. The maize and cowpea were sown in lines, row to row spacing is 45 cm and plant to plant spacing is 25 cm. The seed rate of Maize (20 kg ha<sup>-1</sup>) or Cowpea (18 kg ha<sup>-1</sup>). Maize variety is (1899) or Cowpea (S 263). The data recorded at 30, 60, 90

DAS and at harvest for all the parameters. Harvesting is done after cobs had turned brownish and grain hardened. The harvesting of cowpea was done manually with the help of sickle when pods were matured (when nearly 75% pods had turned brown). Shelling of cobs was done and grains were kept in bags treatment. Threshing was done by beating the bundle with sticks. Seeds were separated by winnowing and kept treatment wise in bags.

### Results and Discussion

#### *Different Growth parameter*

The most decisive factors of growth is plant height (cm), number of leaves plant<sup>-1</sup>, dry weight (g), leaf area index (LAI) have been shown in Table 1 & Fig. 1. The highest value for plant height (172.22 cm), number of leaves plant<sup>-1</sup> (15.55), dry weight (63.29 g), leaf area index (6.14) was recorded with intercropping IC<sub>2</sub> maize + cowpea (2:1) respectively. Significantly maximum values for plant height (174.81 cm), number of leaves plant<sup>-1</sup> (17.53), dry weight (65.03 g), leaf area index (6.44) was recorded with different organic manures and fertilizer levels F<sub>5</sub> (50% RDF + 5 t FYM ha<sup>-1</sup> + 2 t VC ha<sup>-1</sup> + 2 t PM ha<sup>-1</sup>). The probable reason for highest values might due to combined application of organic manures and RDF levels of fertilizers. On the other hand, cowpea is a legume crop having nodules which resulted in higher fixation of atmospheric nitrogen. The legume crop increases the root length and root volume of maize, which helps to absorb more water and nutrients. The similar results were also reported by Murlidhar (2011).

The dry weight plant<sup>-1</sup> was increased due to manures and intercropping with legume improves the physical, chemical as well as biological properties of the soil, which helps in providing suitable environment for growth of plant. The similar results were founded by Asangla and Gohain (2016).

Intercropping can increase the leaf area index and crop growth rate of both crops. A higher leaf area index means that there are more leaves available for photosynthesis, which can lead to higher yields. These findings are in close vicinity of those of Alhaji (2008) and Choudhary *et al.* (2012).

Table 1: Effect of different fertilization on growth parameters of maize with cowpea intercropping.

Treatments	Growth parameters			
	Plant height (cm)	No. of leaves plant <sup>-1</sup>	Dry weight (g)	Leaf Area Index (LAI)
Main plot Intercropping				
IC <sub>1</sub> : Maize + Cowpea (5:1)	168.72	14.51	61.55	5.96
IC <sub>2</sub> : Maize + Cowpea (2:1)	172.22	15.55	63.29	6.14
IC <sub>3</sub> : Maize + Cowpea (3:1)	169.87	14.98	63.23	6.06
SEm(±)	0.21	0.27	0.18	0.01
CD 5%	0.82	1.05	0.71	0.04
Sub plot (Fertilizers)				
F <sub>1</sub> : 100% RDF	167.79	12.55	60.35	5.67
F <sub>2</sub> : 75% RDF +10 t FYM ha <sup>-1</sup>	168.37	13.88	60.79	5.87
F <sub>3</sub> : 50% RDF +5 t FYM ha <sup>-1</sup> + 2t VC ha <sup>-1</sup>	169.76	15.24	63.17	6.05
F <sub>4</sub> : 50% RDF +5t FYM ha <sup>-1</sup> +2 t PM ha <sup>-1</sup>	170.61	15.86	64.14	6.24
F <sub>5</sub> : 50% RDF +5 t FYM ha <sup>-1</sup> +2 t VC ha <sup>-1</sup> +2 t PM ha <sup>-1</sup>	174.81	17.53	65.03	6.44
SEm(±)	1.04	0.47	0.51	0.01
CD 5%	3.00	1.37	1.47	0.02

Table 2: Effect of different fertilization on yield parameters of maize with cowpea intercropping

Treatments	Yield parameters	
	Seed yield (q ha <sup>-1</sup> )	Stover yield (q ha <sup>-1</sup> )
Main plot Intercropping		
IC <sub>1</sub> : Maize + Cowpea (5:1)	42.94	60.39
IC <sub>2</sub> : Maize + Cowpea (2:1)	45.27	66.63
IC <sub>3</sub> : Maize + Cowpea (3:1)	45.02	63.39
SEm(±)	0.07	0.10
CD 5%	0.29	0.40
Sub plot (Fertilizers)		
F <sub>1</sub> : 100% RDF	43.48	61.24
F <sub>2</sub> : 75% RDF +10 t FYM ha <sup>-1</sup>	44.22	62.40
F <sub>3</sub> : 50% RDF +5 t FYM ha <sup>-1</sup> + 2t VC ha <sup>-1</sup>	44.53	62.98
F <sub>4</sub> : 50% RDF +5t FYM ha <sup>-1</sup> +2 t PM ha <sup>-1</sup>	44.74	63.47
F <sub>5</sub> : 50% RDF +5 t FYM ha <sup>-1</sup> +2 t VC ha <sup>-1</sup> +2 t PM ha <sup>-1</sup>	45.09	67.27
SEm(±)	0.11	0.51
CD 5%	0.32	1.48




Fig. 1: Effect of different fertilization on growth parameters of maize with cowpea intercropping




Fig. 2: Effect of different fertilization on yield parameters of maize with cowpea intercropping

The intercropping IC<sub>1</sub> (5:1), IC<sub>2</sub> (2:1), IC<sub>3</sub> (3:1). The fertilizer levels F<sub>1</sub>: 100% RDF, F<sub>2</sub>: 75% RDF +10 t FYM ha<sup>-1</sup>, F<sub>3</sub>: 50% RDF +5 t FYM ha<sup>-1</sup> + 2 t VC ha<sup>-1</sup>, F<sub>4</sub>: 50% RDF +5 t FYM ha<sup>-1</sup> + 2 t PM ha<sup>-1</sup>, F<sub>5</sub>: 50% RDF +5 t FYM ha<sup>-1</sup> + 2 t VC ha<sup>-1</sup> + 2 t PM ha<sup>-1</sup>.

**Yield parameters**

The data on seed yield and stover yield was increased with the organic manure and different levels of fertilizers with intercropping (Table 2 & Fig. 2). The maximum seed yield (45.27 q ha<sup>-1</sup>) and stover yield (66.63 q ha<sup>-1</sup>) were recorded with IC<sub>2</sub> maize + cowpea (2:1) and also seed yield (45.09 q ha<sup>-1</sup>) and stover yield (67.27 q ha<sup>-1</sup>) with F<sub>5</sub> (50% RDF + 5 t FYM ha<sup>-1</sup> + 2 t VC ha<sup>-1</sup> + 2 t PM ha<sup>-1</sup>). The reason behind this proper vegetative and

reproductive growth is better with combined application of fertilizers and manures with intercropping. These also stimulated the various physiological process and metabolic activities which gave better growth and yield of crop. **Conclusion**

On the basis of obtained results from the field experiment, it can be concluded that maximum growth parameters were achieved when the crops intercrop with IC<sub>2</sub> maize + cowpea (2:1) and also with F<sub>5</sub> (50% RDF + 5 t FYM ha<sup>-1</sup> + 2 t VC ha<sup>-1</sup> + 2 t PM ha<sup>-1</sup>).

**References**

Alhaji, I. H. (2008). Yield performance of some cowpea varieties under sole and intercropping with maize at Bauchi Nigeria. *African Research Review*. 2(3): 278 – 291.

- Asangla, H. K. and Gohain, T. (2016). Effect of fodder yield and quality attributes of maize (*Zea mays*) and cowpea (*Vigna unguiculata*) intercropping and different nitrogen levels. *International Journal of Agricultural Science and Research*. 6(2): 349-356.
- Choudhary, V. K.; Suresh Kumar, P. and Bhagawati, R. (2012). Production potential, soil moisture and temperature as influenced by maize- legume intercropping. *International Journal of Science and Nature*. 3(1): 41-46.
- Evans, J.; Mcneil, A. M.; Unkovich, M. J.; Fittell, N. A. and Heena, D. P. (2001). Nat nitrogen balances for cool-season grain legume intercropping and contribution to wheat nitrogen uptake: *A Review. Australian Journal. Exp. Agric.* 41: 347-359.
- Kaviraj, H.; Mansur, C. P.; Vijaymahantesh; Gasti, V. D.; Rajashekhara, E. and Patil, V. (2017). Influence of nutrient management practices on growth and yield of vegetable cowpea in northern dry zone of Karnataka (*Vigna unguiculata* L.). *International Journal of Pure Applied Bioscience*. 5(6): 517-523.
- Li, L.; Tang, C.; Rengel, Z. and Zhang, F. S. (2003). Chickpea facilitates phosphorus uptake by intercropped wheat from an organic phosphorus source. *Journal. Plant Soil* 248: 297-303.
- Mishra, P.; Tiwari, U. S.; Pandey, H. P.; Pathak, R. K. and Sachan, A. K. (2019). Impact of INM on growth and yield of maize (*Zea mays* L.) crop in central plain zone of Uttar Pradesh, India. *International Journal of Current Microbiology and Applied Sciences* 8(4): 138-150.
- Mahmood, F.; Khan, I.; Ashraf, U.; Tanvir, S.; Hussain, S.; Muhammad, S.; Muhammad, A. and Ullah, S. (2017). Effects of organic and inorganic manures on maize and their residual impact on soil physico-chemical properties. *Soil Science and Plant Nutrition*. 17(1): 22-32.
- Murlidhar, D. S. (2011). Production potential of forage maize (*Zea mays* L.) and cowpea (*Vigna unguiculata* L.) intercropping system as influenced by row ratios. Department of Agronomy B.A. College of Agriculture, Anand Agricultural University, Gujarat.
- Reddy, D. M.; Devi, K. B. S. and Sultan, M. A. (2004). Fodder based intercropping for higher biomass and quality fodder production. *Forage Research*. 29(4): 217-218.